



## Primary Series was Pfizer or Moderna

**An additional (3<sup>rd</sup>) dose of mRNA vaccine is recommended for people with moderately or severely weakened immune systems.**

The CDC recommends that moderately to severely immunocompromised people whose primary series was Pfizer or Moderna receive a 3<sup>rd</sup> dose. This includes people who have:

- Been receiving active cancer treatment for tumors or cancers of the blood
- Received an organ transplant and are taking medicine to suppress the immune system
- Received a stem cell transplant within the last 2 years or are taking medicine to suppress the immune system
- Moderate or severe primary immunodeficiency (such as DiGeorge syndrome, Wiskott-Aldrich syndrome)
- Advanced or untreated HIV infection
- Active treatment with high-dose corticosteroids or other drugs that may suppress the immune response


This 3<sup>rd</sup> dose should be the same COVID-19 vaccine as your primary series. Talk to your doctor about the need to get an additional dose of COVID-19 vaccine and about the best timing based on your current treatment plan. This is especially important if you are about to start or restart immunosuppressive treatment.

**A booster dose may be considered at least 6 months after receiving the additional (3<sup>rd</sup>) dose and may be any COVID-19 vaccine.**

### Pfizer – Additional dose should be Pfizer



### Moderna – Additional dose should be Moderna



## Primary Series was Johnson & Johnson (J&J)

The CDC recommends that moderately to severely immunocompromised people whose primary series was J&J receive a booster dose. Talk to your doctor about the best timing based on your current treatment plan. This is especially important if you are about to start or restart immunosuppressive treatment.

